

BRST, anti-BRST and their geometry

L. Bonora^(a,b), R. P. Malik^(c,d)

^(a) *International School for Advanced Studies (SISSA),
Strada Costiera, Via Beirut n.2-4, 34013 Trieste, Italy*

^(b) *INFN, Sezione di Trieste, Trieste, Italy*

^(c) *CAS, Physics Department, BHU, Varanasi-221 005, India*

^(d) *DST-CIMS, Faculty of Science, BHU, Varanasi-221 005, India*

E-mails: bonora@he.sissa.it; malik@bhu.ac.in

Abstract:

We continue the comparison between the field theoretical and geometrical approaches to the gauge field theories of various types, by deriving their Becchi-Rouet-Stora-Tyutin (BRST) and anti-BRST transformation properties and comparing them with the geometrical properties of the bundles and gerbes. In particular, we provide the geometrical interpretation of the so-called Curci-Ferrari conditions that are invoked for the absolute anticommutativity of the BRST and anti-BRST symmetry transformations in the context of non-Abelian 1-form gauge theories as well as Abelian gauge theory that incorporates a 2-form gauge field. We also carry out the explicit construction of the 3-form gauge fields and compare it with the geometry of 2-gerbes.

PACS numbers: 11.15.-q; 03.70.+k

Keywords: Gauge theories, BRST symmetries, gerbes

1 Introduction

In a gauge-fixed quantum gauge field theory, the local gauge symmetry is traded with the nilpotent BRST symmetry. If the gauge-fixing is chosen in a particularly symmetric fashion, the BRST symmetry is accompanied by a twin symmetry, called the anti-BRST. In this paper, we continue the work started in [11] which consists in comparing the language of the quantum field theory with that of the underlying geometry. In particular, in our present work, we concentrate primarily on two topics. The first one is a geometrical interpretation of the BRST and anti-BRST transformations in the twin cases of the non-Abelian 1-form gauge theories and theories incorporating the Abelian 2-form gauge potentials. For the second topic, we focus on the Abelian 3-form gauge field theory and compare it with the 2-gerbe formalism. The above gauge theories, as is well-known, are always endowed with the first-class constraints in the language of Dirac's prescription for the classification scheme [1] [2].

As far as the first topic is concerned, we would like to understand, why in QFT, one needs the constraint relations (referred to as Curci-Ferrari (CF) relations) among the ghost fields and auxiliary fields in order to close the BRST and anti-BRST algebra. We start by reexamining an old problem: in the non-Abelian 1-form gauge field theories, the requirement that the BRST and anti-BRST transformations must anti-commute, imposes a constraint on the (auxiliary) fields of the theory (the so-called CF condition). We will explain, first, how this fact can be given a geometrical interpretation. This type of constraints, however, are characteristic, not only of the non-Abelian 1-form gauge theories, but also of the higher form Abelian (and perhaps non-Abelian) gauge theories whose field contents are based on the concept of gerbes. We wish to give a geometrical interpretation also of these constraints, which we keep referring to as the CF constraints.

Finally, as for our second topic, we introduce a gauge theory based on the Abelian 3-form gauge potential and define its BRST and anti-BRST transformations. Furthermore, we obtain the CF-type conditions for their absolute anti-commutativity. This is to be compared with the geometry of 2-gerbes. To this end we first introduce the latter, find their gauge transformations and show that they can be reduced to the ones determined by the field theoretical methods. Thus, we establish a connection between the field theory of the Abelian 3-form gauge theory with the geometry associated with the 2-gerbes.

2 An old problem: the CF constraints in non-Abelian gauge theories

Let us consider a non-Abelian 1-form gauge theory with gauge group G . We write the gauge potential 1-form as $A = A_\mu^a(x)T^a dx^\mu$ where T^a are the anti-hermitian generators of $Lie(G)$. The appropriate geometry for such a theory is well-known to be a principal fiber bundle $P(M, G)$, the base space being the space-time manifold M and with structure group G . A gauge transformation with infinitesimal parameter $\lambda = \lambda^a(x)T^a$, is given by

$$\delta_\lambda A = D\lambda \equiv d\lambda + [A, \lambda]. \quad (1)$$

Quantizing the theory requires gauge fixing, which, via the Faddeev-Popov procedure, replaces the classical gauge invariance with the BRST invariance. A BRST transformation is analogous to a gauge transformation, except that the gauge parameter is replaced by an anticommuting field $\lambda \rightarrow c = c^a(x)T^a$. The origin of this transmutation was explained in [8]: c represents not a single gauge transformation, but the whole set of gauge transformations. It is, in fact, an alias of the Maurer–Cartan form ω *in the group of gauge transformations*. The fact that ω is a 1-form accounts for the anticommutativity of c . Moreover, contracting it with left invariant vector fields in P gives rise to all the gauge transformations. For this reason, we say that c represents the whole set of gauge transformations. In other words, the ghost field c is the heuristic and compact form that QFT adopts to express the geometric set-up of the non-Abelian 1-form gauge theories.

The BRST transformation (1) is not nilpotent unless we endow c itself with a BRST transformation. Since we have $s^2 A = D(sc) + \frac{1}{2}D([c, c])_+$, the nilpotent BRST transformations must be as follows

$$sA = Dc, \quad sc = -\frac{1}{2}[c, c]_+. \quad (2)$$

Nilpotency is simply a translation, in the language of the QFT, of the fact that the gauge transformations form a Lie algebra. In fact this implies that

$$(\delta_{\lambda_1} \delta_{\lambda_2} - \delta_{\lambda_2} \delta_{\lambda_1})A - \delta_{[\lambda_1, \lambda_2]}A = 0, \quad (3)$$

and we see that (2) exactly mimics this. In particular, the transformation sc mimics the last term at the LHS of (3). This is important for the sequel: *nilpotency of the BRST transformation is the quantum equivalent of the Lie algebra (or Lie group) product law*. In other words, if the BRST transformations were not nilpotent they could not be derived from a classical Lie group product law.

If, in the quantization process, we use the Lorentz gauge fixing, then, the quantum theory turns out to be more symmetric. In this case, we have an additional symmetry, the anti-BRST symmetry. The parameter of this new symmetry is the anti-ghost field $\bar{c} = \bar{c}^a(x)T^a$, which naturally appears in the gauge fixed action.

The complete set of BRST and anti-BRST transformations is [6, 3, 4]

$$\begin{aligned} sA &= Dc, & \bar{s}A &= D\bar{c}, \\ sc &= -\frac{1}{2}[c, c]_+, & \bar{s}\bar{c} &= -\frac{1}{2}[\bar{c}, \bar{c}]_+, \\ s\bar{c} &= B, & \bar{s}c &= \bar{B}, \\ s\bar{B} &= [\bar{B}, c], & \bar{s}B &= [B, \bar{c}], \end{aligned} \quad (4)$$

the remaining transformations being trivial. The fields $B = B_\mu^a(x)T^a dx^\mu$ and $\bar{B} = \bar{B}_\mu^a(x)T^a dx^\mu$ are auxiliary scalar commuting fields. It is easy to see that s and $\bar{s}$ are nilpotent. One expects the total BRST-anti-BRST operator $s + \bar{s}$ to be nilpotent too. We have

$$(s\bar{s} + \bar{s}s)A = D(B + \bar{B} + [c, \bar{c}]_+). \quad (5)$$

Therefore nilpotency of $s + \bar{s}$ requires that

$$B + \bar{B} + [c, \bar{c}]_+ = 0. \quad (6)$$

This is the original CF constraint. On the other hand, we have seen above that nilpotency corresponds to Lie algebra structure. Therefore, it is natural to require that (6) hold good. The nilpotency of $s + \bar{s}$ (which corresponds to anticommutativity) has been used in the literature to study the physical spectrum of non-Abelian 1-form gauge theories (see, e.g. [9]). Added to it, the requirement of nilpotency of BRST and anti-BRST transformations is, since long, a standard folklore in the realm of BRST formalism.

Let us now delve into the geometrical interpretation of $\bar{s}$. The field $\bar{c}$ is a sort of copy of c . Therefore, it would seem that the natural geometrical setting for a BRST and anti-BRST-symmetric theory is $P(M, G \times G)$, that is a principal fiber bundle with structure group $G \times G$, with c taking values in the Lie algebra of the first group and $\bar{c}$ in the Lie algebra of the second. But this cannot be the case since $[c, \bar{c}] \neq 0$.

We propose the following geometrical set-up. Starting from two isomorphic principal fiber bundles $P(M, G)$ and $Q(M, G')$ on the same base space, we can easily construct the bundle $(P + Q)(M, G \times G')$ on M with structure group $G \times G'$, by pulling back the product $P(M, G) \times Q(M, G')$ via the diagonal map $\Delta(x) = (x, x)$. The fibers of $P + Q$ are couples of fibers of P and Q , i.e. they are (p, q) such that $\pi_P(p) = x = \pi_Q(q)$. The transition functions

$\hat{\psi}_{\alpha\beta}$ of $P + Q$ are given by

$$\hat{\psi}_{\alpha\beta} \equiv (\psi_{\alpha\beta}, \psi'_{\alpha\beta}) : U_\alpha \cap U_\beta \rightarrow G \times G', \quad (7)$$

where $\psi_{\alpha\beta}$ and $\psi'_{\alpha\beta}$ are the transition functions of P and Q , respectively.

Now, if $G \equiv G'$, in $\hat{\psi}_{\alpha\beta}$ we can choose, in particular, $\psi_{\alpha\beta} \equiv \psi'_{\alpha\beta}$. This means that $P + Q$ is reducible to a bundle $R(M, G_d)$, with structure group $G_d = \text{Diag}(G \times G)$ (see [10], Prop.5.3). It is obvious that the fibers of R are the diagonal fibers of $P + Q$. R represents the geometric set-up we are looking for. It is isomorphic to $P(M, G)$, as it should be, but what matters is the fine structure of this isomorphism. In fact, it hosts two gauge groups, two copies of G , and this is what we need to accommodate the above cited two different types of gauge transformations.

Let us consider the relation between R and $P + Q$. The reduction is specified by a homomorphism $R \rightarrow P + Q$, defined by a function f such that $\forall u \in R$, $f(u) \in P + Q$ which reduces to the identity on the base M , and a group homomorphism $f_\delta : G \rightarrow \text{Diag}(G \times G)$, such that $f(ug) = f(u)f_\delta(g)$ for any $u \in R$ and $g \in G$. A connection $\mathcal{A}$ in $P + Q$ reduces to a connection A in R and the two are related by, [10],

$$f^* \mathcal{A} = f_\delta \cdot A, \quad (8)$$

where f^* denotes the pull-back and $(f_\delta \cdot A)(X) = df_\delta(A(X)) \equiv (A(X), A(X))$ for any vector field X in R . The correspondence between A and $\mathcal{A}$ is one-to-one (see, e.g. [10] ch.II.6).

Let us consider now finite gauge transformations. They are given by vertical automorphisms ψ of a principal bundle, i.e., by bundle morphisms that do not affect the basis. In the case of R , the fibers get transformed as $\psi((p, p)) = (\psi(p), \psi(p))$, where ψ is an automorphism of $P = Q$. Now let us consider the same construction of $P + Q$ as above, but with P replaced by $\psi_1^* P$, where ψ_1 is now an automorphism of P (as opposed to an automorphism of Q), i.e., we consider $\psi_1^* P + Q$ where $\psi_1^* P$ is the gauge transformed form of P . Naturally we would get a version of R with fibers $(\psi_1(p), q)$, where $q = \psi_1(q)$. Therefore, trivially $\psi_1((p, q)) = (\psi_1(p), q)$, which defines an automorphism ψ_1 of R originating from an automorphism of P . In the same way, we can get automorphisms ψ_2 originated from the automorphisms of Q .

Taking a connection A in R , we get, therefore, two types of gauge transformations $\psi_1^* A$ and $\psi_2^* A$. We link them to c and $\bar{c}$, respectively. In this way c ($\bar{c}$) is associated to the Maurer-Cartan form of the first (second) factor in $G \times G$, respectively. But they are both projected to the diagonal group. As a consequence, the anticommutator $[c, \bar{c}]_+$ is non-vanishing. This seems to be the most appropriate geometrical set-up for the BRST and anti-BRST.

Next, let us consider two infinitesimal gauge transformations of the first type (ψ_1) and call them λ_1 and λ_2 , and two of the second type $\bar{\lambda}_1$ and $\bar{\lambda}_2$. It is easy to prove that

$$\left((\delta_{\lambda_1} + \delta_{\bar{\lambda}_1})(\delta_{\lambda_2} + \delta_{\bar{\lambda}_2}) - (\delta_{\lambda_2} + \delta_{\bar{\lambda}_2})(\delta_{\lambda_1} + \delta_{\bar{\lambda}_1}) \right) A - \delta_{[\lambda_1 + \bar{\lambda}_1, \lambda_2 + \bar{\lambda}_2]} A = 0. \quad (9)$$

This is the geometrical meaning of (4). However, in the anticommuting language, we cannot reproduce it without introducing the auxiliary fields B and $\bar{B}$, for we have

$$(s\bar{s} + \bar{s}s)A = D(s\bar{c} + \bar{s}c + [c, \bar{c}]_+), \quad (10)$$

which motivates the definition of B and $\bar{B}$ in (4).

We stress again that (4) and (6) express, in the language of quantum field theory, the simple Lie algebra rule (9).

Finally, let us make a comment concerning the Abelian case. When the gauge group G is Abelian, we can, of course, repeat everything word by word. In equations (3) and (4) the Lie brackets vanish, so BRST and anti-BRST transformations are disconnected. The CF constraint is replaced by $B + \bar{B} = 0$. These auxiliary fields are, in fact, decoupled and one can do without them. We would like to point out, however, that the 1-form Abelian bundle case is the only one in which the CF constraint is superfluous. In the next more complicated case of the Abelian 2-form, which corresponds to the geometry of 1-gerbes, the CF constraint is essential.

3 CF constraints for 1-gerbes

The BRST and anti-BRST transformations for the 1-gerbes were worked out in [11]. Here we would like to give the relevant geometrical interpretation.

Let us recall some basic definitions. A 1-gerbe (for mathematical properties of gerbes see [12, 13], for previous applications in physics see [15]) may be characterized by a triple (B, A, f) , formed by the 2-form B , 1-form A and 0-form f , respectively. These are related in the following way. Given a covering $\{U_i\}$ of the manifold M , we associate to each U_i a two-form B_i . On a double intersection $U_i \cap U_j$, we have $B_i - B_j = dA_{ij}$. On the triple intersections $U_i \cap U_j \cap U_k$, we must have $A_{ij} + A_{jk} + A_{ki} = df_{ijk}$. Finally, on the quadruple intersections $U_i \cap U_j \cap U_k \cap U_l$, the following integral cocycle condition must be satisfied:

$$f_{ijl} - f_{ijk} + f_{jkl} - f_{ikl} = 2\pi n, \quad n = 0, 1, 2, 3, \dots \quad (11)$$

This integrality condition will not concern us in our Lagrangian formulation but it has to be imposed as an external condition.

Two triples, represented by (B, A, f) and (B', A', f') , respectively, are gauge equivalent if they satisfy the following relations

$$B'_i = B_i + dC_i \quad \text{on } U_i, \quad (12)$$

$$A'_{ij} = A_{ij} + C_i - C_j + d\lambda_{ij} \quad \text{on } U_i \cap U_j, \quad (13)$$

$$f'_{ijk} = f_{ijk} + \lambda_{ij} + \lambda_{ki} + \lambda_{jk} \quad \text{on } U_i \cap U_j \cap U_k, \quad (14)$$

for the one-forms C and the zero-forms λ .

We will now define the BRST and anti-BRST transformations corresponding to these geometrical transformations. As shown in [11], one can proceed in two different ways. Either one defines an action for the triple of local fields (B, A, f) , quantizes it by adding all the ghost and auxiliary fields that are needed, and verifies that the quantum action has the two BRST and anti-BRST symmetries below (this is what we do in section 4 for the 3-form gauge field). Or, more heuristically, by analogy with the 1-form gauge theory, one starts from the (known) gauge transformations of the (B, A, f) fields and constructs the BRST and anti-BRST transformations by simply relying on nilpotency and consistency. The two procedures lead to the same results up to minor ambiguities (see below). It should be recalled that while the above geometric transformations are defined on (multiple) neighborhood overlaps, the BRST and anti-BRST transformations in quantum field theory are defined on a single local coordinate patch. These (local, field-dependent) transformations are the means QFT uses to record the underlying geometry.

The appropriate BRST and anti-BRST transformations are¹

$$\begin{aligned} s B &= dC, & s A &= C + d\lambda, & s f &= \lambda + \mu, \\ s C &= -d\beta, & s \lambda &= \beta, & s \mu &= -\beta, \\ s \bar{C} &= -K, & s \bar{K} &= d\rho, & s \bar{\mu} &= -g, \\ s \bar{\beta} &= -\bar{\rho}, & s \bar{\lambda} &= g, & s \bar{g} &= \rho, \end{aligned} \quad (15)$$

together with $s [\rho, \bar{\rho}, g, K_\mu, \beta] = 0$, and

$$\begin{aligned} \bar{s} B &= d\bar{C}, & \bar{s} A &= \bar{C} + d\bar{\lambda}, & \bar{s} f &= \bar{\lambda} + \bar{\mu}, \\ \bar{s} \bar{C} &= +d\bar{\beta}, & \bar{s} \bar{\lambda} &= -\bar{\beta}, & \bar{s} \bar{\mu} &= -\bar{\beta}, \\ \bar{s} C &= +\bar{K}, & \bar{s} K &= -d\bar{\rho}, & \bar{s} \bar{\mu} &= \bar{g}, \\ \bar{s} \beta &= +\rho, & \bar{s} \lambda &= -\bar{g}, & \bar{s} g &= -\bar{\rho}, \end{aligned} \quad (16)$$

while $\bar{s} [\bar{\beta}, \bar{g}, \bar{K}_\mu, \rho, \bar{\rho}] = 0$.

¹As done in [11], we take into account here also the scalar field f .

It can be easily verified that $(s + \bar{s})^2 = 0$ if the following constraint is satisfied:

$$\bar{K} - K = d\bar{g} - dg. \quad (17)$$

This condition is both BRST and anti-BRST invariant. It is the analogue of the Curci-Ferrari condition in non-Abelian 1-form gauge theories and we will refer to it with the same name.

The field content and BRST and anti-BRST structure for 1-gerbe field theories is shown schematically in Figure 1.




Figure 1: *A schematic view of BRST and anti-BRST transformations for 1-gerbes*

Before we proceed with the discussion, we would like to note that the above realization of the BRST and anti-BRST algebra is not the only possibility. In general, it may be possible to augment it by the addition of a sub-algebra of elements which are all in the kernel of both s and $\bar{s}$ or, if it contains such a sub-algebra, the latter could be moded out. For instance, in equations (15,16) ρ and $\bar{\rho}$ form an example of this type of subalgebra. It is easy to see that ρ and $\bar{\rho}$ can be consistently set equal to 0.

We would like now to suggest a geometrical setting for these transformations and the relevant CF constraints. We have already noted that for Abelian gauge bundles, we can carry out a construction similar to that of the previous section. The Lie bracket terms in (4) are trivial and we can, in fact, dispense with the auxiliary fields B and $\bar{B}$. That is to say the BRST and anti-BRST transformations are independent of any interference term and the CF condition is unnecessary. The gauge transformations, underlying

a 1-gerbe, are of the Abelian type too, but, contrary to the 1-form gauge theory (i.e. 0-gerbe), they produce a non-trivial CF condition (17).

The geometric set-up for the BRST and anti-BRST 1-gerbe transformations is similar to the one that is true for the gauge bundles. One can define both Cartesian products and pullbacks of 1-gerbes, (see, e.g. [13]). We can, therefore, take the two copies of the same 1-gerbe and make their product. Then, we pull the result back by the diagonal map (see above). So far, the construction is parallel to the one in the previous section. Now we have to do the analogue of the mapping to the diagonal of $G \times G$. This is not so easy in the above formulation of the 1-gerbes. But there is another formulation due to Hitchin [14] which we now recall and dwell upon. With respect to a covering $\{U_i\}$, a 1-gerbe is specified by the following data:

- a line bundle L_{ij} for any intersection $U_i \cap U_j$,
- an isomorphism between L_{ij} and L_{ji}^{-1} ,
- a trivialization of $L_{ij}L_{jk}L_{ki}$, that is a map $\theta_{ijk} : U_i \cap U_j \cap U_k \longrightarrow U(1)$,
- this map satisfies the cocycle condition $\theta_{jkl}\theta_{ikl}^{-1}\theta_{ijl}\theta_{ijk}^{-1} = 1$.

In this definition, the line bundles are associated with the $U(1)$ principal bundles, L^{-1} represents the bundle dual to L , and the products of the bundles are tensor products. It is now possible to redo the construction of the previous section (including the diagonal mapping for fibers) locally, i.e., for each line bundle and each map θ (by replacing each local P with a corresponding isomorphic R) while satisfying the conditions of the 1-gerbe definition. This means that we can think of two copies of the transformations (12–14) with parameters C_i, λ_{ij} and $\bar{C}_i, \bar{\lambda}_{ij}$, respectively.

This is the classical set-up we propose in order to accommodate the BRST and anti-BRST 1-gerbe transformations.

From the two copies of (12–14), it is easy to recognize the origin of many of the transformations in (15,16). For instance from (12), we see that C_i is not uniquely defined, we could choose $C'_i = C_i + d\beta_i$ (or $\bar{C}'_i = \bar{C}_i + d\bar{\beta}_i$), obtaining in this way, the classical analogue of sC (or $\bar{s}\bar{C}$). Plugging this into (13), we get $\lambda'_{ij} = \lambda_{ij} - \beta_i + \beta_j$ and obtain the analogue of $s\lambda$, etc. Denoting now, for simplicity, the classical infinitesimal transformations δ and $\bar{\delta}$ corresponding to the two copies of (12–14), we get the following

$$0 = (\delta\bar{\delta} - \bar{\delta}\delta)B = \delta(d\bar{C}) - \bar{\delta}(dC) = d(\delta\bar{C} - \bar{\delta}C), \quad (18)$$

which is nothing but the geometric origin of the CF constraint.

4 Abelian 3–form gauge theories and 2–gerbes

In this section, we would like to give a more complex example, that of the Abelian 3–form gauge theory, which has not been explicitly worked out as yet. In fact, we derive the proper BRST and anti–BRST symmetry transformations within a field theoretical approach. In the next section, we will compare it with the geometric setting of 2–gerbes.

The nilpotent and absolutely anticommuting (anti-)BRST symmetry transformations for the Abelian 3–form theory have been derived [7] by exploiting the geometrical superfield formalism [5]. Section 4.1 is devoted to the discussion of the (anti-)BRST invariance of the coupled Lagrangian densities of the theory. Our Sec.4.2 deals with the discrete and ghost scale symmetry transformations for the ghost part of the Lagrangian densities. We obtain the algebraic structures, satisfied by the conserved charges (corresponding to the above continuous symmetries), in Sec. 4.3.

4.1 Preliminary: nilpotent (anti-)BRST symmetries in superfield formulation

By exploiting the superfield approach to BRST formalism [5], the nilpotent and absolutely anticommuting (anti-)BRST symmetry transformations for the free Abelian 3–form gauge theory have been derived in [7]. To be more precise, such a theory² can be embedded in a $(D, 2)$ -dimensional supermanifold and the application of the celebrated horizontality condition leads to the following (anti-)BRST symmetry transformations³

$$\begin{aligned} sB_{\mu\nu\eta} &= \partial_\mu C_{\nu\eta} + \partial_\nu C_{\eta\mu} + \partial_\eta C_{\mu\nu}, & sC_{\mu\nu} &= \partial_\mu \beta_\nu - \partial_\nu \beta_\mu, \\ s\bar{C}_{\mu\nu} &= B_{\mu\nu}, & s\bar{B}_{\mu\nu} &= \partial_\mu f_\nu - \partial_\nu f_\mu, & s\bar{\beta}_\mu &= \bar{F}_\mu, & s\beta_\mu &= \partial_\mu C_2, \\ sF_\mu &= -\partial_\mu B, & s\bar{f}_\mu &= \partial_\mu B_1, & s\bar{C}_2 &= B_2, & sC_1 &= -B, \\ s\phi_\mu &= f_\mu, & s\bar{C}_1 &= B_1, & s[C_2, f_\mu, \bar{F}_\mu, B, B_1, B_2, B_{\mu\nu}] &= 0, \end{aligned} \quad (19)$$

$$\begin{aligned} \bar{s}B_{\mu\nu\eta} &= \partial_\mu \bar{C}_{\nu\eta} + \partial_\nu \bar{C}_{\eta\mu} + \partial_\eta \bar{C}_{\mu\nu}, & \bar{s}\bar{C}_{\mu\nu} &= \partial_\mu \bar{\beta}_\nu - \partial_\nu \bar{\beta}_\mu, \\ \bar{s}C_{\mu\nu} &= \bar{B}_{\mu\nu}, & \bar{s}B_{\mu\nu} &= \partial_\mu \bar{f}_\nu - \partial_\nu \bar{f}_\mu, & \bar{s}\beta_\mu &= F_\mu, & \bar{s}\bar{\beta}_\mu &= \partial_\mu \bar{C}_2, \end{aligned}$$

²In [7], the theory was considered in 4D, but the results hold in a D dimensional Minkowski space-time. We take here the Greek indices $\mu, \nu, \eta, \dots = 0, 1, \dots, D-1$ to correspond to the spacetime directions of the flat Minkowski spacetime manifold endowed with a metric with signatures $(+1, -1, \dots, -1)$. For algebraic convenience, we have changed $\bar{B} \rightarrow B, B_{\mu\nu}^{(2)} \rightarrow B_{\mu\nu}, \bar{B}_{\mu\nu}^{(1)} \rightarrow \bar{B}_{\mu\nu}$ and exchanged $F_\mu \leftrightarrow \bar{F}_\mu$ in the notations of [7].

³In a field theoretic approach, it is more convenient to use the component notation rather than the synthetic differential form language.

$$\begin{aligned}\bar{s}\bar{F}_\mu &= -\partial_\mu B_2, & \bar{s}f_\mu &= -\partial_\mu B_1, & \bar{s}C_2 &= B, & \bar{s}C_1 &= -B_1, \\ \bar{s}\phi_\mu &= \bar{f}_\mu, & \bar{s}\bar{C}_1 &= -B_2, & \bar{s}[\bar{C}_2, \bar{f}_\mu, F_\mu, B, B_1, B_2, \bar{B}_{\mu\nu}] &= 0,\end{aligned}\quad (20)$$

where $B_{\mu\nu\eta}$ is the totally antisymmetric tensor gauge field, $(\bar{C}_{\mu\nu})C_{\mu\nu}$ are the fermionic antisymmetric (anti-)ghost fields with ghost number $(-1)+1$, $(\bar{\beta}_\mu)\beta_\mu$ are the Lorentz vector bosonic ghost-for-ghost (anti-)ghost fields with ghost number $(-2)+2$ and $(\bar{C}_2)C_2$ are the fermionic ghost-for-ghost-for-ghost (anti-)ghost Lorentz scalar fields with ghost number $(-3)+3$. The vector field ϕ_μ and auxiliary fields B, B_1, B_2 are bosonic in nature and fermionic fields $(f_\mu)f_\mu$ and $(\bar{F}_\mu)F_\mu$ are the auxiliary (anti-)ghost fields with ghost numbers $(-1)+1$, respectively. Furthermore, it will be noted that the bosonic auxiliary fields B and B_2 carry ghost number $+2$ and -2 , respectively.

The above off-shell nilpotent ($s_{(a)b}^2 = 0$) (anti-)BRST symmetry transformations are not absolutely anticommuting in nature because

$$\{s, \bar{s}\}B_{\mu\nu\eta} \neq 0, \quad \{s, \bar{s}\}C_{\mu\nu} \neq 0, \quad \{s, \bar{s}\}\bar{C}_{\mu\nu} \neq 0. \quad (21)$$

It is interesting, however, to mention that the superfield formalism [7] yields the following CF type restrictions

$$\begin{aligned}f_\mu + F_\mu &= \partial_\mu C_1, & \bar{f}_\mu + \bar{F}_\mu &= \partial_\mu \bar{C}_1, \\ B_{\mu\nu} + \bar{B}_{\mu\nu} &= \partial_\mu \phi_\nu - \partial_\nu \phi_\mu,\end{aligned}\quad (22)$$

which ensure the absolute anticommutativity (i.e. $s\bar{s} + \bar{s}s \equiv \{s, \bar{s}\} = 0$) of the above nilpotent (anti-)BRST symmetry transformations.

We wrap up this section with the following comments:

(i) the nilpotency and absolute anticommutativity of the (anti-)BRST symmetry transformations are the indispensable consequences of the superfield approach to BRST formalism [5],

(ii) there are *three* CF type restriction (cf. (21)) for the 4D Abelian 3-form gauge theory whereas there is one each for the 4D Abelian 2-form [11] and 4D (non-)Abelian 1-form gauge theories [6],

(iii) two of the CF type restrictions in (21) are fermionic in nature and one of them is bosonic (whereas for the Abelian 2-form and (non-)Abelian 1-form gauge theories only bosonic type CF restriction exists). For the Abelian 1-form gauge theory, the CF type restriction is *trivial*,

(iv) the CF type restriction is one of the key properties of any arbitrary p -form gauge theory described in the framework of BRST formalism, and

(v) the CF type restrictions, that emerge from superfield formalism, are always (anti-)BRST invariant relationships (e.g. $(s, \bar{s}) [f_\mu + F_\mu - \partial_\mu C_1] = 0$, $(s, \bar{s}) [\bar{f}_\mu + \bar{F}_\mu - \partial_\mu \bar{C}_1] = 0$, $(s, \bar{s}) [B_{\mu\nu} + \bar{B}_{\mu\nu} - (\partial_\mu \phi_\nu - \partial_\nu \phi_\mu)] = 0$).

4.2 Lagrangian densities: nilpotent (anti-)BRST symmetry transformations

With the help of the (anti-)BRST symmetry transformations (18) and (19), one can derive the (anti-)BRST invariant Lagrangian density for the Abelian 3-form gauge theory as

$$\begin{aligned}
\mathcal{L}_B &= \frac{1}{24} H^{\mu\nu\eta\xi} H_{\mu\nu\eta\xi} + s\bar{s} \left[\frac{1}{2} \bar{C}_2 C_2 - \frac{1}{2} \bar{C}_1 C_1 - \frac{1}{2} \bar{C}_{\mu\nu} C^{\mu\nu} - \bar{\beta}_\mu \beta^\mu \right. \\
&\quad \left. - \frac{1}{2} \phi_\mu \phi^\mu - \frac{1}{6} B_{\mu\nu\eta} B^{\mu\nu\eta} \right], \\
\mathcal{L}_{\bar{B}} &= \frac{1}{24} H^{\mu\nu\eta\xi} H_{\mu\nu\eta\xi} - \bar{s}s \left[\frac{1}{2} \bar{C}_2 C_2 - \frac{1}{2} \bar{C}_1 C_1 - \frac{1}{2} \bar{C}_{\mu\nu} C^{\mu\nu} - \bar{\beta}_\mu \beta^\mu \right. \\
&\quad \left. - \frac{1}{2} \phi_\mu \phi^\mu - \frac{1}{6} B_{\mu\nu\eta} B^{\mu\nu\eta} \right], \tag{23}
\end{aligned}$$

where $H_{\mu\nu\eta\xi} = \partial_\mu B_{\nu\eta\xi} - \partial_\nu B_{\eta\xi\mu} + \partial_\eta B_{\xi\mu\nu} - \partial_\xi B_{\mu\nu\eta}$ is the totally antisymmetric curvature tensor derived from the 4-form $H^{(4)} = dB^{(3)} \equiv [(dx^\mu \wedge dx^\nu \wedge dx^\eta \wedge dx^\xi)]/[(4)!] H_{\mu\nu\eta\xi}$. Here $d = dx^\mu \partial_\mu$ (with $d^2 = 0$) is the exterior derivative and the 3-form $B^{(3)} = [(dx^\mu \wedge dx^\nu \wedge dx^\eta)]/[(3)!] B_{\mu\nu\eta}$ defines the totally antisymmetric tensor gauge field $B_{\mu\nu\eta}$. It will be noted that, within the square brackets of (22), we have taken the combination of terms that are Lorentz scalars, carry the ghost number equal to zero and, furthermore, they possess the proper mass dimensions.

Modulo some total spacetime derivatives, the explicit computation of the square bracketed terms yield the following:

$$\begin{aligned}
&s\bar{s} \left[\frac{1}{2} \bar{C}_2 C_2 - \frac{1}{2} \bar{C}_1 C_1 - \frac{1}{2} \bar{C}_{\mu\nu} C^{\mu\nu} - \bar{\beta}_\mu \beta^\mu - \frac{1}{2} \phi_\mu \phi^\mu - \frac{1}{6} B_{\mu\nu\eta} B^{\mu\nu\eta} \right] \\
&= (\partial_\mu B^{\mu\nu\eta}) B_{\nu\eta} + \frac{1}{2} B_{\mu\nu} \bar{B}^{\mu\nu} + (\partial_\mu \bar{C}_{\nu\eta} + \partial_\nu \bar{C}_{\eta\mu} + \partial_\eta \bar{C}_{\mu\nu}) (\partial^\mu C^{\nu\eta}) \\
&\quad - (\partial_\mu \bar{\beta}_\nu - \partial_\nu \bar{\beta}_\mu) (\partial^\mu \beta^\nu) - B B_2 - \frac{1}{2} B_1^2 + (\partial_\mu \bar{C}^{\mu\nu}) f_\nu - (\partial_\mu C^{\mu\nu}) \bar{F}_\nu \\
&\quad + \partial_\mu \bar{C}_2 \partial^\mu C_2 + \bar{f}^\mu f_\mu - \bar{F}^\mu F_\mu + (\partial \cdot \beta) B_2 + (\partial \cdot \phi) B_1 - (\partial \cdot \bar{\beta}) B, \tag{24}
\end{aligned}$$

$$\begin{aligned}
&-\bar{s}s \left[\frac{1}{2} \bar{C}_2 C_2 - \frac{1}{2} \bar{C}_1 C_1 - \frac{1}{2} \bar{C}_{\mu\nu} C^{\mu\nu} - \bar{\beta}_\mu \beta^\mu - \frac{1}{2} \phi_\mu \phi^\mu - \frac{1}{6} B_{\mu\nu\eta} B^{\mu\nu\eta} \right] \\
&= -(\partial_\mu B^{\mu\nu\eta}) \bar{B}_{\nu\eta} + \frac{1}{2} B_{\mu\nu} \bar{B}^{\mu\nu} + (\partial_\mu \bar{C}_{\nu\eta} + \partial_\nu \bar{C}_{\eta\mu} + \partial_\eta \bar{C}_{\mu\nu}) (\partial^\mu C^{\nu\eta}) \\
&\quad - (\partial_\mu \bar{\beta}_\nu - \partial_\nu \bar{\beta}_\mu) (\partial^\mu \beta^\nu) - B B_2 - \frac{1}{2} B_1^2 - (\partial_\mu \bar{C}^{\mu\nu}) F_\nu + (\partial_\mu C^{\mu\nu}) \bar{f}_\nu \\
&\quad + \partial_\mu \bar{C}_2 \partial^\mu C_2 + \bar{f}^\mu f_\mu - \bar{F}^\mu F_\mu + (\partial \cdot \beta) B_2 + (\partial \cdot \phi) B_1 - (\partial \cdot \bar{\beta}) B. \tag{25}
\end{aligned}$$

The difference in the above explicit computations is due to the fact that the (anti-)BRST symmetry transformations are anticommuting only on the constrained surface defined by the CF type conditions (21).

With the help of the CF type restrictions (21), we can re-express the Lagrangian density $\mathcal{L}_B$ in an appropriate form as⁴

$$\begin{aligned}\mathcal{L}_B = & \frac{1}{24}H^{\mu\nu\eta\xi}H_{\mu\nu\eta\xi} + B^{\mu\nu}\left(\partial^\eta B_{\eta\mu\nu} + \frac{1}{2}[\partial_\mu\phi_\nu - \partial_\nu\phi_\mu]\right) \\ & - \frac{1}{2}B_{\mu\nu}B^{\mu\nu} + (\partial_\mu\bar{C}_{\nu\eta} + \partial_\nu\bar{C}_{\eta\mu} + \partial_\eta B_{\mu\nu})(\partial^\mu C^{\nu\eta}) - (\partial \cdot \bar{\beta})B \\ & - (\partial_\mu\bar{\beta}_\nu - \partial_\nu\bar{\beta}_\mu)(\partial^\mu\beta^\nu) - BB_2 - \frac{1}{2}B_1^2 + (\partial_\mu\bar{C}^{\mu\nu} + \partial^\nu\bar{C}_1)f_\nu \\ & - (\partial_\mu C^{\mu\nu} - \partial^\nu C_1)\bar{F}_\nu + \partial_\mu\bar{C}_2\partial^\mu C_2 + (\partial \cdot \beta)B_2 + (\partial \cdot \phi)B_1.\end{aligned}\quad (26)$$

It should be noted that, in the above, we have taken the expression (23) but have replaced $\bar{B}_{\mu\nu}, \bar{f}_\mu, F_\mu$ by exploiting the CF type conditions in (21). In an exactly similar fashion, the Lagrangian density $\mathcal{L}_{\bar{B}}$ can be re-written as

$$\begin{aligned}\mathcal{L}_{\bar{B}} = & \frac{1}{24}H^{\mu\nu\eta\xi}H_{\mu\nu\eta\xi} - \bar{B}^{\mu\nu}\left(\partial^\eta B_{\eta\mu\nu} - \frac{1}{2}[\partial_\mu\phi_\nu - \partial_\nu\phi_\mu]\right) \\ & - \frac{1}{2}\bar{B}_{\mu\nu}\bar{B}^{\mu\nu} + (\partial_\mu\bar{C}_{\nu\eta} + \partial_\nu\bar{C}_{\eta\mu} + \partial_\eta B_{\mu\nu})(\partial^\mu C^{\nu\eta}) - (\partial \cdot \bar{\beta})B \\ & - (\partial_\mu\bar{\beta}_\nu - \partial_\nu\bar{\beta}_\mu)(\partial^\mu\beta^\nu) - BB_2 - \frac{1}{2}B_1^2 - (\partial_\mu\bar{C}^{\mu\nu} + \partial^\nu\bar{C}_1)F_\nu \\ & + (\partial_\mu C^{\mu\nu} - \partial^\nu C_1)\bar{f}_\nu + \partial_\mu\bar{C}_2\partial^\mu C_2 + (\partial \cdot \beta)B_2 + (\partial \cdot \phi)B_1.\end{aligned}\quad (27)$$

In the above, it should be noted that we have taken the expression (24) but have substituted for $B_{\mu\nu}, f_\mu, \bar{F}_\mu$ by exploiting the CF type conditions in (21).

Under the BRST symmetry transformations (18), the above Lagrangian densities transform as

$$\begin{aligned}s\mathcal{L}_B = & \partial_\mu\left[(\partial^\mu C^{\nu\eta} + \partial^\nu C^{\eta\mu} + \partial^\eta C^{\mu\nu})B_{\nu\eta} + B^{\mu\nu}f_\nu\right. \\ & \left. - (\partial^\mu\beta^\nu - \partial^\nu\beta^\mu)\bar{F}_\nu + B_1f^\mu - B\bar{F}^\mu + B_2\partial^\mu C_2\right],\end{aligned}\quad (28)$$

$$\begin{aligned}s\mathcal{L}_{\bar{B}} = & -\partial_\mu\left[(\partial^\mu C^{\nu\eta} + \partial^\nu C^{\eta\mu} + \partial^\eta C^{\mu\nu})\bar{B}_{\nu\eta} + B^{\mu\nu}F_\nu\right. \\ & \left. - (\partial^\mu\beta^\nu - \partial^\nu\beta^\mu)\bar{f}_\nu - B_1f^\mu + B\bar{F}^\mu - B_2\partial^\mu C_2\right. \\ & \left. + B^{\mu\nu\eta}(\partial_\nu f_\eta - \partial_\eta f_\nu) + \bar{C}^{\mu\nu}\partial_\nu B + C^{\mu\nu}\partial_\nu B_1\right] + X,\end{aligned}\quad (29)$$

⁴In fact, there are, in total, six more possibilities to express the equivalent Lagrangian densities with the help of CF type restrictions (21). It is, however, only the equivalent forms like (25) and (26) that have perfect transformations like (27) and (28).

where the extra piece X , in the above, is given below

$$\begin{aligned}
X = & (\partial^\mu C^{\nu\eta} + \partial^\nu C^{\eta\mu} + \partial^\eta C^{\mu\nu}) \partial_\mu [\bar{B}_{\nu\eta} + B_{\nu\eta} - (\partial_\nu \phi_\eta - \partial_\eta \phi_\nu)] \\
& + B^{\mu\nu} \partial_\mu [f_\nu + F_\nu - \partial_\nu C_1] - [\bar{B}_{\mu\nu} + B_{\mu\nu} - (\partial_\mu \phi_\nu - \partial_\nu \phi_\mu)] (\partial^\mu f^\nu) \\
& + (\bar{f}^\mu + \bar{F}^\mu - \partial^\mu \bar{C}_1) (\partial_\mu B) - (f^\mu + F^\mu - \partial^\mu C_1) (\partial_\mu B_1) \\
& - (\partial^\mu \bar{\beta}^\nu - \partial^\nu \bar{\beta}^\mu) \partial_\mu [(\bar{f}_\nu + \bar{F}_\nu - \partial_\nu \bar{C}_1)].
\end{aligned} \tag{30}$$

Thus, we note that both the above coupled Lagrangian densities are equivalent with respect to the BRST transformations (18) on the submanifold defined by the CF type field equations (21). In other words, if we impose the CF type restrictions, both the above Lagrangian densities respect the BRST symmetry transformations because they transform to total derivatives.

In exactly the above fashion, we note that, under the anti-BRST symmetry transformations (19), the coupled Lagrangian densities transform as

$$\begin{aligned}
\bar{s}\mathcal{L}_{\bar{B}} = & \partial_\mu \left[(\partial^\mu \bar{C}^{\nu\eta} + \partial^\nu \bar{C}^{\eta\mu} + \partial^\eta \bar{C}^{\mu\nu}) \bar{B}_{\nu\eta} + \bar{B}^{\mu\nu} \bar{f}_\nu \right. \\
& \left. - (\partial^\mu \bar{\beta}^\nu - \partial^\nu \bar{\beta}^\mu) F_\nu + B_1 \bar{f}^\mu + B_2 F^\mu - B \partial^\mu \bar{C}_2 \right],
\end{aligned} \tag{31}$$

$$\begin{aligned}
\bar{s}\mathcal{L}_B = & \partial_\mu \left[(\partial^\mu \bar{C}^{\nu\eta} + \partial^\nu \bar{C}^{\eta\mu} + \partial^\eta \bar{C}^{\mu\nu}) B_{\nu\eta} - \bar{B}^{\mu\nu} \bar{F}_\nu \right. \\
& + (\partial^\mu \bar{\beta}^\nu - \partial^\nu \bar{\beta}^\mu) f_\nu + B_1 \bar{f}^\mu + B_2 F^\mu - B \partial^\mu \bar{C}_2 \\
& \left. + (B^{\mu\nu\eta}) (\partial_\nu \bar{f}_\eta - \partial_\eta \bar{f}_\nu) + \bar{C}^{\mu\nu} \partial_\nu B_1 - C^{\mu\nu} \partial_\nu B_2 \right] + Y,
\end{aligned} \tag{32}$$

where the extra piece Y , in the above equation, is

$$\begin{aligned}
Y = & -(\partial^\mu \bar{C}^{\nu\eta} + \partial^\nu \bar{C}^{\eta\mu} + \partial^\eta \bar{C}^{\mu\nu}) \partial_\mu [\bar{B}_{\nu\eta} + B_{\nu\eta} - (\partial_\nu \phi_\eta - \partial_\eta \phi_\nu)] \\
& + \bar{B}^{\mu\nu} \partial_\mu [\bar{f}_\nu + \bar{F}_\nu - \partial_\nu \bar{C}_1] - [\bar{B}_{\mu\nu} + B_{\mu\nu} - (\partial_\mu \phi_\nu - \partial_\nu \phi_\mu)] (\partial^\mu \bar{f}^\nu) \\
& - (\bar{f}^\mu + \bar{F}^\mu - \partial^\mu \bar{C}_1) (\partial_\mu B_1) - (f^\mu + F^\mu - \partial^\mu C_1) (\partial_\mu B_2) \\
& - (\partial^\mu \bar{\beta}^\nu - \partial^\nu \bar{\beta}^\mu) \partial_\mu [(f_\nu + F_\nu - \partial_\nu C_1)].
\end{aligned} \tag{33}$$

It is gratifying to note that both the above coupled Lagrangian densities respect the off-shell nilpotent (anti-)BRST symmetry transformations on the submanifold of the spacetime that is described by the constrained CF type field conditions (21). These features, under the (anti-)BRST symmetry transformations, are exactly like the ones we come across in the context of the BRST approach to 4D non-Abelian 1-form gauge theory (see, e.g. [9]).

4.3 Ghost Lagrangian density: global scale and discrete symmetry transformations

The ghost parts of the Lagrangian densities (25) and (26), even though they look quite different in their appearance, are actually *equal* on the submanifold defined by the CF type restrictions (21). Thus, let us take one of them, as

$$\begin{aligned}\mathcal{L}_{(g)}^{(B)} &= (\partial_\mu \bar{C}_{\nu\eta} + \partial_\nu \bar{C}_{\eta\mu} + \partial_\eta \bar{C}_{\mu\nu})(\partial^\mu C^{\nu\eta}) - (\partial \cdot \bar{\beta})B \\ &- (\partial_\mu \bar{\beta}_\nu - \partial_\nu \bar{\beta}_\mu)(\partial^\mu \beta^\nu) - BB_2 + (\partial_\mu \bar{C}^{\mu\nu} + \partial^\nu \bar{C}_1)f_\nu \\ &- (\partial_\mu C^{\mu\nu} - \partial^\nu C_1)\bar{F}_\nu + \partial_\mu \bar{C}_2 \partial^\mu C_2 + (\partial \cdot \beta)B_2.\end{aligned}\quad (34)$$

The above Lagrangian density respects the following continuous global scale transformations for the ghost fields, namely;

$$\begin{aligned}C_{\mu\nu} &\rightarrow e^{+\Omega}C_{\mu\nu}, & \bar{C}_{\mu\nu} &\rightarrow e^{-\Omega}\bar{C}_{\mu\nu}, & C_1 &\rightarrow e^{+\Omega}C_1, \\ \bar{C}_1 &\rightarrow e^{-\Omega}\bar{C}_1, & f_\mu &\rightarrow e^{+\Omega}f_\mu, & F_\mu &\rightarrow e^{+\Omega}F_\mu, \\ \bar{f}_\mu &\rightarrow e^{-\Omega}\bar{f}_\mu, & \bar{F}_\mu &\rightarrow e^{-\Omega}\bar{F}_\mu, & \beta_\mu &\rightarrow e^{+2\Omega}\beta_\mu, \\ \bar{\beta}_\mu &\rightarrow e^{-2\Omega}\bar{\beta}_\mu, & B &\rightarrow e^{+2\Omega}B, & B_2 &\rightarrow e^{-2\Omega}B_2, \\ C_2 &\rightarrow e^{+3\Omega}C_2, & \bar{C}_2 &\rightarrow e^{-3\Omega}\bar{C}_2,\end{aligned}\quad (35)$$

where Ω is a global scale infinitesimal parameter and the numbers, present in the exponentials, denote the ghost numbers of the corresponding dynamical and/or auxiliary (anti-)ghost fields.

In addition to the above global continuous symmetry transformations, the above Lagrangian density $\mathcal{L}_g^{(B)}$ also respects the following discrete symmetry transformations amongst the (anti-)ghost fields, namely;

$$\begin{aligned}C_{\mu\nu} &\rightarrow \pm i \bar{C}_{\mu\nu}, & C_1 &\rightarrow \mp i \bar{C}_1, & \beta_\mu &\rightarrow \pm i \bar{\beta}_\mu, \\ \bar{C}_{\mu\nu} &\rightarrow \pm i C_{\mu\nu}, & \bar{C}_1 &\rightarrow \mp i C_1, & \bar{\beta}_\mu &\rightarrow \mp i \beta_\mu, \\ f_\mu &\rightarrow \pm i \bar{F}_\mu, & B_2 &\rightarrow \pm i B, & C_2 &\rightarrow \pm i \bar{C}_2, \\ \bar{F}_\mu &\rightarrow \pm i f_\mu, & B &\rightarrow \mp i B_2, & \bar{C}_2 &\rightarrow \pm i C_2.\end{aligned}\quad (36)$$

The above symmetry transformations enable us to go from the BRST symmetry transformations to the anti-BRST symmetry transformations and *vice-versa*. The above transformations have been written for the ghost part of the Lagrangian density of $\mathcal{L}_B$. However, similar kind of transformations can be written out for the ghost part of the Lagrangian density of $\mathcal{L}_{\bar{B}}$ where, in addition to the above transformations, we shall require

$$\bar{f}_\mu \rightarrow \pm i F_\mu, \quad F_\mu \rightarrow \pm i \bar{f}_\mu. \quad (37)$$

We close this section with the remark that the ghost part of the coupled Lagrangian densities remain invariant under (34), (35) and (36). Furthermore, it can be checked that the CF type restrictions also remain invariant under the transformations (34), (35) and (36) because $f_\mu + F_\mu = \pm \partial_\mu C_1$ and $\bar{f}_\mu + \bar{F}_\mu = \pm \partial_\mu \bar{C}_1$ are allowed by the anticommutativity property. It is trivial to state that $B_{\mu\nu\eta} \rightarrow B_{\mu\nu\eta}$, $B_{\mu\nu} \rightarrow B_{\mu\nu}$, $\bar{B}_{\mu\nu} \rightarrow \bar{B}_{\mu\nu}$, $\phi_\mu \rightarrow \phi_\mu$ under the ghost transformations because these fields carry ghost number equal to zero.

4.4 Conserved charges: algebraic structures

It is clear from equations (27) and (28) that the Lagrangian densities $\mathcal{L}_B$ and $\mathcal{L}_{\bar{B}}$ transform precisely to the total spacetime derivative under the BRST and anti-BRST transformations (18) and (19). According to the Noether's theorem, we have the following expressions for the conserved currents

$$\begin{aligned} J_{(B)}^\mu &= s\Phi_i \frac{\partial \mathcal{L}_B}{\partial(\partial_\mu \Phi_i)} - Z^\mu, \\ J_{(\bar{B})}^\mu &= \bar{s}\Phi_i \frac{\partial \mathcal{L}_{\bar{B}}}{\partial(\partial_\mu \Phi_i)} - S^\mu, \end{aligned} \quad (38)$$

where the generic dynamical field $\Phi_i = B_{\mu\nu\eta}, C_{\mu\nu}, \bar{C}_{\mu\nu}, \beta_\mu, \bar{\beta}_\mu, \phi_\mu, \bar{C}_1, C_1, \bar{C}_2, \bar{C}_2$ and the explicit expression for Z^μ and S^μ are (cf. (27) and (30))

$$\begin{aligned} Z^\mu &= (\partial^\mu C^{\nu\eta} + \partial^\nu C^{\eta\mu} + \partial^\eta C^{\mu\nu}) B_{\nu\eta} + B^{\mu\nu} f_\nu \\ &\quad - (\partial^\mu \beta^\nu - \partial^\nu \beta^\mu) \bar{F}_\nu + B_1 f^\mu - B \bar{F}^\mu + B_2 \partial^\mu C_2, \\ S^\mu &= (\partial^\mu \bar{C}^{\nu\eta} + \partial^\nu \bar{C}^{\eta\mu} + \partial^\eta \bar{C}^{\mu\nu}) \bar{B}_{\nu\eta} + \bar{B}^{\mu\nu} \bar{f}_\nu \\ &\quad - (\partial^\mu \bar{\beta}^\nu - \partial^\nu \bar{\beta}^\mu) F_\nu + B_1 \bar{f}^\mu + B_2 F^\mu - B \partial^\mu \bar{C}_2. \end{aligned} \quad (39)$$

The conserved currents that emerge from the above equations are as follows

$$\begin{aligned} J_{(B)}^\mu &= H^{\mu\nu\eta\xi} (\partial_\nu C_{\eta\xi}) + (\partial^\mu C^{\nu\eta} + \partial^\nu C^{\eta\mu} + \partial^\eta C^{\mu\nu}) B_{\nu\eta} + B^{\mu\nu} f_\nu + B_1 f^\mu \\ &\quad + B_2 \partial^\mu C_2 - B \bar{F}^\mu - (\partial^\mu \bar{\beta}^\nu - \partial^\nu \bar{\beta}^\mu) (\partial_\nu C_2) - (\partial^\mu \beta^\nu - \partial^\nu \beta^\mu) \bar{F}_\nu \\ &\quad - (\partial^\mu \bar{C}^{\nu\eta} + \partial^\nu \bar{C}^{\eta\mu} + \partial^\eta \bar{C}^{\mu\nu}) (\partial_\nu \beta_\eta - \partial_\eta \beta_\nu), \end{aligned} \quad (40)$$

$$\begin{aligned} J_{(\bar{B})}^\mu &= H^{\mu\nu\eta\xi} (\partial_\nu \bar{C}_{\eta\xi}) + (\partial^\mu \bar{C}^{\nu\eta} + \partial^\nu \bar{C}^{\eta\mu} + \partial^\eta \bar{C}^{\mu\nu}) \bar{B}_{\nu\eta} + \bar{B}^{\mu\nu} \bar{f}_\nu + B_1 \bar{f}^\mu \\ &\quad - B \partial^\mu \bar{C}_2 + B_2 F^\mu - (\partial^\mu \beta^\nu - \partial^\nu \beta^\mu) (\partial_\nu \bar{C}_2) - (\partial^\mu \bar{\beta}^\nu - \partial^\nu \bar{\beta}^\mu) F_\nu \\ &\quad + (\partial^\mu C^{\nu\eta} + \partial^\nu C^{\eta\mu} + \partial^\eta C^{\mu\nu}) (\partial_\nu \bar{\beta}_\eta - \partial_\eta \bar{\beta}_\nu), \end{aligned} \quad (41)$$

where all the derivatives of (37) have been calculated from $\mathcal{L}_B$ and $\mathcal{L}_{\bar{B}}$.

The conservation law (i.e. $\partial_\mu J_{(i)}^\mu = 0, i = B, \bar{B}$) can be proven by taking into account the following equations of motion that are derived from $\mathcal{L}_B$

$$\begin{aligned}
& \square B_{\mu\nu\eta} = 0, \quad B_{\mu\nu} = (\partial^\eta B_{\eta\mu\nu}) + \frac{1}{2}(\partial_\mu \phi_\nu - \partial_\nu \phi_\mu), \quad \square \beta_\mu = 0, \\
& \square \bar{\beta}_\mu = 0, \quad \partial \cdot F = 0, \quad \partial \cdot \bar{F} = 0, \quad \partial \cdot f = 0, \quad \partial_\mu B^{\mu\nu} + \partial^\nu B_1 = 0, \\
& \partial \cdot \bar{f} = 0, \quad \partial_\mu C^{\mu\nu} = \partial^\nu C_1, \quad \partial_\mu \bar{C}^{\mu\nu} = -\partial^\nu \bar{C}_1, \\
& \square C_1 = 0, \quad \square \bar{C}_1 = 0, \quad \square C_2 = 0, \quad \square \bar{C}_2 = 0, \quad \square B_1 = 0, \\
& B = \partial \cdot \beta, \quad B_1 = \partial \cdot \phi, \quad B_2 = -(\partial \cdot \bar{\beta}), \quad \square \phi_\mu + \partial_\mu (\partial \cdot \phi) = 0, \\
& \square C_{\mu\nu} + \frac{1}{2}(\partial_\mu f_\nu - \partial_\nu f_\mu) = 0, \quad \square \bar{C}_{\mu\nu} + \frac{1}{2}(\partial_\mu \bar{f}_\nu - \partial_\nu \bar{f}_\mu) = 0, \\
& \partial_\mu H^{\mu\nu\eta\xi} + (\partial^\nu B^{\eta\xi} + \partial^\eta B^{\xi\nu} + \partial^\xi B^{\nu\eta}) = 0.
\end{aligned} \tag{42}$$

The Euler-Lagrange equations of motion, that emerge from $\mathcal{L}_{\bar{B}}$, are the same as the above but for the following differences, namely;

$$\begin{aligned}
& \bar{B}_{\mu\nu} = -(\partial^\eta B_{\eta\mu\nu}) + \frac{1}{2}(\partial_\mu \phi_\nu - \partial_\nu \phi_\mu), \quad \partial_\mu \bar{B}^{\mu\nu} + \partial^\nu B_1 = 0, \\
& \square C_{\mu\nu} - \frac{1}{2}(\partial_\mu f_\nu - \partial_\nu f_\mu) = 0, \quad \square \bar{C}_{\mu\nu} - \frac{1}{2}(\partial_\mu \bar{f}_\nu - \partial_\nu \bar{f}_\mu) = 0, \\
& \partial_\mu H^{\mu\nu\eta\xi} - (\partial^\nu \bar{B}^{\eta\xi} + \partial^\eta \bar{B}^{\xi\nu} + \partial^\xi \bar{B}^{\nu\eta}) = 0.
\end{aligned} \tag{43}$$

It is elementary to state that (41) and (42) are exploited, in a judicious manner, for the proof of the conservation laws.

The continuous ghost symmetry transformations of (34) lead to the derivation of the ghost conserved current as given below

$$\begin{aligned}
J_{(g)}^\mu &= (\partial^\mu \bar{C}^{\nu\eta} + \partial^\nu \bar{C}^{\eta\mu} + \partial^\eta \bar{C}^{\mu\nu}) C_{\nu\eta} + (\partial^\mu C^{\nu\eta} + \partial^\nu C^{\eta\mu} + \partial^\eta C^{\mu\nu}) \bar{C}_{\nu\eta} \\
&- 2(\partial^\mu \bar{\beta}^\nu - \partial^\nu \bar{\beta}^\mu) \beta_\nu + 2(\partial^\mu \beta^\nu - \partial^\nu \beta^\mu) \bar{\beta}_\nu + 2B \bar{\beta}^\mu + 2B_2 \beta^\mu \\
&+ 3(\partial^\mu \bar{C}_2) C_2 - 3\bar{C}_2 \partial^\mu C_2 - \bar{C}^{\mu\nu} f_\nu - C^{\mu\nu} \bar{f}_\nu + C_1 \bar{F}^\mu - \bar{C}_1 f^\mu.
\end{aligned} \tag{44}$$

The conservation law (i.e. $\partial_\mu J_{(g)}^\mu = 0$) can be proven by taking the help of equations of motion (41) and (42). The conserved (i.e. $\dot{Q}_{(r)} = 0, r = B, \bar{B}, g$) charges (i.e. $Q_{(r)} = \int d^3x J_{(r)}^0, r = B, \bar{B}, g$), that emerge from the conserved currents (39), (40) and (43), are as follows

$$\begin{aligned}
Q_{(B)} &= \int d^3x \left[H^{0ijk} (\partial_i C_{jk}) + (\partial^0 C^{\nu\eta} + \partial^\nu C^{\eta 0} + \partial^\eta C^{0\nu}) B_{\nu\eta} + B_1 f^0 \right. \\
&- (\partial^0 \bar{\beta}^i - \partial^i \bar{\beta}^0) \partial_i C_2 - (\partial^0 \beta^i - \partial^i \beta^0) \bar{F}_i + B^{0i} f_i + B_2 \dot{C}_2 \\
&- \left. (\partial^0 \bar{C}^{\nu\eta} + \partial^\nu \bar{C}^{\eta 0} + \partial^\eta \bar{C}^{0\nu}) (\partial_\nu \beta_\eta - \partial_\eta \beta_\nu) - B \bar{F}^0 \right],
\end{aligned} \tag{45}$$

$$\begin{aligned}
Q_{(\bar{B})} = & \int d^3x \left[H^{0ijk} (\partial_i \bar{C}_{jk}) - (\partial^0 \bar{C}^{\nu\eta} + \partial^\nu \bar{C}^{\eta 0} + \partial^\eta \bar{C}^{0\nu}) \bar{B}_{\nu\eta} + B_1 \bar{f}^0 \right. \\
& - (\partial^0 \beta^i - \partial^i \beta^0) \partial_i \bar{C}_2 - (\partial^0 \bar{\beta}^i - \partial^i \bar{\beta}^0) F_i + \bar{B}^{0i} \bar{f}_i - B \dot{\bar{C}}_2 \\
& \left. + (\partial^0 C^{\nu\eta} + \partial^\nu C^{\eta 0} + \partial^\eta C^{0\nu}) (\partial_\nu \bar{\beta}_\eta - \partial_\eta \bar{\beta}_\nu) + B_2 F^0 \right], \quad (46)
\end{aligned}$$

$$\begin{aligned}
Q_{(g)} = & \int d^3x \left[3 \dot{\bar{C}}_2 C_2 - 3 \bar{C}_2 \dot{C}_2 + (\partial^0 \bar{C}^{\nu\eta} + \partial^\nu \bar{C}^{\eta 0} + \partial^\eta \bar{C}^{0\nu}) C_{\nu\eta} - \bar{C}_1 f^0 \right. \\
& + 2(\partial^0 \beta^i - \partial^i \beta^0) \bar{\beta}_i - 2(\partial^0 \bar{\beta}^i - \partial^i \bar{\beta}^0) \beta_i - \bar{C}^{0i} f_i - C^{0i} \bar{F}_i \\
& \left. + 2B \bar{\beta}^0 + 2B_2 \beta^0 + C_1 \bar{F}^0 + (\partial^0 C^{\nu\eta} + \partial^\nu C^{\eta 0} + \partial^\eta C^{0\nu}) \bar{C}_{\nu\eta} \right]. \quad (47)
\end{aligned}$$

The above conserved charges are the generators of the nilpotent and continuous (anti-)BRST as well as continuous ghost scale transformations.

The application of the continuous symmetry transformations on the above charges produces the following algebra

$$\begin{aligned}
sQ_{(B)} = -i\{Q_{(B)}, Q_{(B)}\} = 0 & \Rightarrow Q_{(B)}^2 = 0, \\
\bar{s}Q_{(\bar{B})} = -i\{Q_{(\bar{B})}, Q_{(\bar{B})}\} = 0 & \Rightarrow Q_{(\bar{B})}^2 = 0, \\
sQ_{(\bar{B})} = -i\{Q_{(\bar{B})}, Q_{(B)}\} = 0 & \Rightarrow \{Q_{(\bar{B})}, Q_{(B)}\} = 0, \\
sQ_{(g)} = -i[Q_{(g)}, Q_{(B)}] = -Q_{(B)} & \Rightarrow i[Q_{(g)} Q_{(B)}] = +Q_{(B)}, \\
\bar{s}Q_{(g)} = -i[Q_{(g)}, Q_{(\bar{B})}] = +Q_{(\bar{B})} & \Rightarrow i[Q_{(g)} Q_{(\bar{B})}] = -Q_{(\bar{B})}. \quad (48)
\end{aligned}$$

The above algebra is the standard algebra in the BRST formalism. In the above, we have not written some more transformations on charges. However, those are implied from the above (e.g. $\bar{s}Q_{(B)} = \{Q_{(B)}, Q_{(\bar{B})}\} = 0$, etc.). The above algebra ensures that the ghost number of the BRST charge is (+1) and that of the anti-BRST charge is (-1).

5 BRST and anti-BRST for a 2-gerbe field theory

A 2-gerbe can be described by a quadruple (C, B, A, f) , defined by a 3-form C , a 2-form B , a 1-form A and a scalar f with the following relations. Given a covering $\{U_i\}$ of M , we associate to each U_i a three-form C_i . On a double intersection $U_i \cap U_j$, we have $C_i - C_j = dB_{ij}$. On the triple intersections $U_i \cap U_j \cap U_k$, we must have $B_{ij} + B_{jk} + B_{ki} = dA_{ijk}$. On quadruple intersections $U_i \cap U_j \cap U_k \cap U_l$, we must have $A_{ijl} - A_{ijk} + A_{jkl} - A_{ikl} = df_{ijkl}$ and on quintuple

intersection $U_i \cap U_j \cap U_k \cap U_l \cap U_m$ the following integral cocycle condition must be satisfied:

$$f_{ijlm} - f_{ijkm} + f_{ijkl} - f_{iklm} + f_{jklm} = 2\pi n, \quad n = 0, 1, 2, 3, \dots \quad (49)$$

Two quadruples (C, B, A, f) and (C', B', A', f') are gauge equivalent if they satisfy

$$B'_{ij} = B_{ij} + \Gamma_i - \Gamma_j + db_{ij}, \quad (50)$$

$$A'_{ijk} = A_{ijk} + b_{ij} + b_{jk} + b_{ki} + d\gamma_{ijk}, \quad (51)$$

$$f'_{ijkl} = f_{ijkl} + \gamma_{ijl} - \gamma_{ijk} + \gamma_{jkl} - \gamma_{ikl}. \quad (52)$$




Figure 2: A *skematic view of BRST and anti-BRST transformations for 2-gerbes*

Starting from these transformation properties, we can define the BRST and anti-BRST transformations appropriate for the field contents of a 2-gerbe field theory. We have to introduce a whole lot of auxiliary fields, even more than in the previous cases, in order to capture its ample gauge freedom

in the language of the absolute anticommutativity property. The overall field content is summarized in Figure 2.

The BRST and anti-BRST transformations, for the boundary faces of the figure, are as follows

$$\begin{aligned}
sC &= d\Gamma, & s\Gamma &= db, & sb &= d\epsilon, \\
sB &= \Gamma + d\gamma, & s\gamma &= -b + dt, & st &= \epsilon = st', \\
sA &= \gamma + \eta + d\xi, & s\eta &= b - dt', & s\xi &= t - t' = -s\zeta, \\
sf &= \xi + \zeta,
\end{aligned} \tag{53}$$

and

$$\begin{aligned}
\bar{s}C &= d\bar{\Gamma}, & \bar{s}\bar{\Gamma} &= d\bar{b}, & \bar{s}\bar{b} &= d\bar{\epsilon}, \\
\bar{s}B &= \bar{\Gamma} + d\bar{\gamma}, & \bar{s}\bar{\gamma} &= -\bar{b} + d\bar{t}, & \bar{s}\bar{t} &= \bar{\epsilon} = \bar{s}\bar{t}', \\
\bar{s}A &= \bar{\gamma} + \bar{\eta} + d\bar{\xi}, & \bar{s}\bar{\eta} &= \bar{b} - d\bar{t}', & \bar{s}\bar{\xi} &= \bar{t} - \bar{t}' = -\bar{s}\bar{\zeta}, \\
\bar{s}f &= \bar{\xi} + \bar{\zeta},
\end{aligned} \tag{54}$$

The remaining transformations are

$$\begin{aligned}
s\bar{\Gamma} &= K, & s\bar{b} &= -\bar{\rho}, & s\bar{K} &= d\rho, \\
s\bar{\gamma} &= \bar{X} + d\bar{g}, & s\bar{\eta} &= -\bar{X} - d\bar{g}, & sg &= -\lambda, \\
sX &= -\rho, & s\bar{X} &= -d\lambda, & s\bar{g} &= \lambda, \\
s\bar{\xi} &= s\bar{\zeta} = \frac{1}{2}(g + \bar{g}), & s\bar{t} &= s\bar{t}' = -\bar{\lambda},
\end{aligned} \tag{55}$$

and

$$\begin{aligned}
\bar{s}\Gamma &= \bar{K}, & \bar{s}b &= -\rho, & \bar{s}K &= d\bar{\rho}, \\
\bar{s}\gamma &= X + dg, & \bar{s}\eta &= -X - dg, & \bar{s}g &= -\bar{\lambda}, \\
\bar{s}\bar{X} &= -\bar{\rho}, & \bar{s}X &= d\bar{\lambda}, & \bar{s}\bar{g} &= \bar{\lambda}, \\
\bar{s}\xi &= \bar{s}\zeta = -\frac{1}{2}(g + \bar{g}), & \bar{s}t &= \bar{s}t' = \lambda.
\end{aligned} \tag{56}$$

Moreover, we also have $s[\epsilon, \bar{\epsilon}, K, \rho, \bar{\rho}, \lambda, \bar{\lambda}] = 0$, and $\bar{s}[\epsilon, \bar{\epsilon}, \bar{K}, \rho, \bar{\rho}, \lambda, \bar{\lambda}] = 0$,

With these transformation properties, the anticommutator of the BRST and anti-BRST transformations ($s\bar{s} + \bar{s}s$) annihilates all the above fields, except for C and A , which require the CF condition

$$K + \bar{K} + d(X + \bar{X}) = 0. \tag{57}$$

Once again, we believe, observing the above BRST and anti-BRST algebra, that a geometric set-up, similar to the one for 1-gerbes, could be constructed. However, since a mathematical formulation (analogous to Hitchin's for 1-gerbes) is still missing for 2-gerbes, we leave this problem open.

5.1 Correspondence with the field theory model

Let us now compare the formulas of this section with the field-theoretical derivation of sec. 4. It is clear that in sec. 4 we considered a reduced model, where the only non-ghost field is the three-form field. The correspondence with that Abelian 3-form gauge theory can be seen as follows. Suppress the two-form, one-form and 0-form (non-ghost) fields of the previous subsection. For the remaining fields the correspondence is as follows:

$$\begin{array}{lll}
B_{\mu\nu\lambda} \leftrightarrow C, & C_{\mu\nu} \leftrightarrow \Gamma & \bar{C}_{\mu\nu} \leftrightarrow \bar{\Gamma}, \\
\beta_\mu \leftrightarrow b, & \bar{\beta}_\mu \leftrightarrow \bar{b}, & B_{\mu\nu} \leftrightarrow K, \\
\bar{B}_{\mu\nu} \leftrightarrow \bar{K}, & f_\mu \leftrightarrow \rho, & F_\mu \leftrightarrow -\rho, \\
\bar{f}_\mu \leftrightarrow \bar{\rho}, & \bar{F}_\mu \leftrightarrow -\bar{\rho}, & C_2 \leftrightarrow \epsilon, \\
\bar{C}_2 \leftrightarrow \bar{\epsilon}, & C_1 \leftrightarrow \lambda, & \bar{C}_1 \leftrightarrow \bar{\lambda}, \\
B \leftrightarrow 0, & B_1 \leftrightarrow 0, & B_2 \leftrightarrow 0
\end{array} \tag{58}$$

and

$$\phi_\mu \leftrightarrow -X - \bar{X} \tag{59}$$

The ghost fields B, B_1, B_2 are an example of subalgebra that can be moded out. Upon moding it out, the first two CF constraints in equation (21) turn out to be trivial.

6 Conclusion

In this paper, we have proposed a geometrical interpretation of the BRST and anti-BRST algebra in an ordinary non-Abelian 1-form gauge field theory. To be more precise, we have provided a geometrical interpretation of the CF constraints, which are needed in order to close the algebra. For this purpose, we have slightly modified the well-known geometric setting of the gauge theories based on the principal fiber bundles to allow for the non-trivial anti-commutator between ghosts and anti-ghosts fields.

We have remarked that if the gauge group is Abelian, the CF conditions are pointless. If an Abelian structure, however, involves a 2-form gauge field, the non-trivial CF constraints turn up, once again. The geometry of such theories is that of 1-gerbes. This geometry dictates the BRST and anti-BRST structure separately, but a modification is necessary in order to interpret the absolute anticommutativity of the joint BRST and anti-BRST transformations. Based on the previous example of 1-form gauge field theories, we have suggested how this modification can be implemented.

Finally, we have studied the case of Abelian gauge field theories involving 3-form fields. We have derived the relevant BRST and anti-BRST transformations in the framework of quantum field theory (essentially by using the superfield method) and, in particular, we have worked out the specific CF conditions. The geometry relevant to this kind of theories is based on the 2-gerbe structure. We have derived the BRST and anti-BRST transformations from the 2-gerbe geometry and shown that they are compatible with those obtained via field theory methods. The geometrical interpretation of the CF conditions, in this context, has still to be spelled out in detail.

Acknowledgements

One of us (RPM) would like to acknowledge the financial support from DST, Govt. of India, under the SERC project grant No: SR/S2/HEP-23/2006. He would also like to express his deep sense of gratitude to the Director, AS-ICTP, for the warm hospitality where a part of this work was done.

References

- [1] P. A. M. Dirac, *Lectures on quantum mechanics*, Belfer Graduate School of Science, Yeshiva University Press, New York, 1964.
- [2] K. Sundermeyer, *Constrained dynamics: Lecture notes in physics*, vol. 169, Springer-Verlag, Berlin, 1982.
- [3] S. Weinberg, *The quantum theory of fields: Modern applications*, vol. 2, Cambridge University Press, Cambridge, 1996.
- [4] M. Henneaux, C. Teitelboim, *Quantization of fields with constraints*, Princeton University Press, Princeton (NJ), 1992.
- [5] See, e.g., L. Bonora and M. Tonin, *Superfield formulation of extended BRS symmetry*, Phys. Lett. **98** (1981) 48.
- [6] G. Curci and R. Ferrari, *Slavnov transformations and supersymmetry*, Phys. Lett. **B 63** (1976) 91.
- [7] R. P. Malik, *Abelian 2-form gauge theory: superfield formalism*, Eur. Phys. J. C **60** (2009) 457, arXiv:0702.0039 [hep-th].

- [8] L.Bonora, P.Cotta-Ramusino, *Some Remarks on BRS Transformations, Anomalies and the Cohomology of the Lie Algebra of the Group of Gauge Transformations*, Comm.Math.Phys. **87** (1983) 589.
- [9] N. Nakanishi and I. Ojima, *Covariant operator formalism of gauge theories and quantum gravity*, World Scientific, Singapore 1990.
- [10] S.Kobayashi and K.Nomizu, *Foundations of Differential Geometry* vol.I, John Wiley & Sons, New York, 1963.
- [11] L. Bonora and R. P. Malik, *BSRT, anti-BRST and gerbes*, Phys. Lett. B **655** (2007) 75 [arXiv:0707.3922 [hep-th]].
- [12] J. Giraud, *Cohomologie non-abélienne*, Grundle. **179**, Springer Verlag, Berlin (1971).
- [13] J.-L. Brylinski, *Characteristic Classes and Geometric Quantization*, Progr. in Math. **107**, Birkhäuser, Boston (1993).
- [14] N. Hitchin, *Lectures on Special Lagrangian Submanifolds*, arXiv:math/9907034
- [15] A. L. Carey, J. Mickelsson and M. K. Murray, *Bundle gerbes applied to quantum field theory*, Rev. Math. Phys. **12** (2000) 65 [arXiv:hep-th/9711133].
J. Kalkkinen, *Gerbes and massive type II configurations*, JHEP **9907** (1999) 002 [arXiv:hep-th/9905018].
P. Bouwknegt, A. L. Carey, V. Mathai, M. K. Murray and D. Stevenson, *Twisted K-theory and K-theory of bundle gerbes*, Commun. Math. Phys. **228** (2002) 17 [arXiv:hep-th/0106194].
M. I. Caicedo, I. Martin and A. Restuccia, *Gerbes and duality*, Annals Phys. **300** (2002) 32 [arXiv:hep-th/0205002].
K. Gawedzki and N. Reis, *WZW branes and gerbes*, Rev. Math. Phys. **14** (2002) 1281 [arXiv:hep-th/0205233].
P. Aschieri, L. Cantini and B. Jurco, *Nonabelian bundle gerbes, their differential geometry and gauge theory*, Commun. Math. Phys. **254** (2005) 367 [arXiv:hep-th/0312154].
J. M. Isidro, *Gerbes and Heisenberg's uncertainty principle*, Int. J. Geom. Meth. Mod. Phys. **3** (2006) 1469 [arXiv:hep-th/0512241].